## **Brandon J. Barnes, M.D.** | Orthopaedic Surgeon, Sports Medicine Specialist BrandonBarnesMD.com (239) 334-7000

## Meniscal Allograft Transplantation (MAT) Protocol

	Weight-baring	Brace	Range of Motion	Exercises
Phase 1 0-2 weeks	Heel touch only	Locked in full extension at all times	Gentle passive 0- 90°	Calf pumps, patellar mobilization, quadriceps tightening sets and SLR in brace
		Off for hygiene and home exercise only		
Phase 2	<b>2-6 weeks:</b> Heel touch only	<b>2-6 weeks:</b> Unlocked 0-90°	Advance as tolerated	<b>2-6 weeks:</b> Add side-lying hip and core, advance quad set and stretching (a)
2-8 weeks	<b>6-8 weeks:</b> Advance 25% weekly and progress to full with normalized gait pattern	Discontinue brace at 6 weeks	CPM continues for 6 hours per day 0-90°	
				<b>6-8 weeks:</b> Addition of heel raises, total gym (closed chain), gait normalization, eccentric quads, eccentric hamstrings
				Advance core work with hip, gluteal and core strengthening
				No weight-bearing with flexion >90°
Phase 3	Full	None	Full	Progress closed chain activities
8-12 weeks				Advance hamstring exercises, proprioception & balance exercises, lunges & leg press 0- 90° only
				Advance core work with hip, gluteal and core strengthening
				Begin Stationary Bike at 10 weeks
Phase 4	Full	None	Full	Progress Phase III exercises and
3-5 months				<b>functional activities</b> : walking lunges, planks, bridges, Swiss ball, half-Bosu exercises

				Advance core work with hip, gluteal and core strengthening			
				Swimming allowed at 16 weeks			
Phase 5	Full	None	Full	Advance non-impact activity			
5-12				No running, jumping, pivoting, or sports			
months				participation unless cleared by MD			
CPM – continuous passive motion machine; SLR – Straight-leg raise							
(a) Avoid tibial rotation for first 8 weeks.							